•LERUP SSP

HLA-C*04:09N Product Insert Page 1 of 8
101.861-12 – including *Taq* polymerase, IFU-01 Visit https://labproducts.caredx.com for 101.861-12u – without *Taq* polymerase, IFU-02 "Instructions for Use" (IFU)

Lot No.: **4L4** Lot-specific information

Olerup SSP® HLA-C*04:09N

Product number: 101.861-12 – including *Taq* polymerase

101.861-12u – without *Taq* polymerase

Lot number: 4L4

Expiry date: 2024-08-01

Number of tests: 12 Number of wells per test: 2+1

Storage - pre-aliquoted primers: dark at -20°C

PCR Master Mix: -20°C
 Adhesive PCR seals
 Product Insert
 RT

This Product Description is only valid for Lot No. 4L4.

Complete product documentation consists of generic Instructions for Use (IFU), lot specific Product Insert, Worksheet and Certificate.

The HLA-C*04:09N is unchanged compared to the previous *Olerup* SSP® HLA-C*04:09N Lot (5F7).

The HLA-C*04:09N specificity and interpretation tables has been updated for the HLA-C alleles described since the previous *Olerup* SSP® HLA-C*04:09N lot **(Lot No. 5F7)** was made. The kit design is based on IMGT/HLA database 3.40.0.

"Instructions for Use" (IFU)

Page 2 of 8

Lot No.: 4L4 Lot-specific information

Well 3 contains Negative Control primer pairs, that will amplify the majority of the Olerup SSP® HLA Class I, DRB, DQB1, DPB1 and DQA1 amplicons as well as all the amplicons generated by the control primer pairs matching the human growth hormone gene.

HLA-specific PCR product sizes range from 75 to 200 base pairs. The PCR product generated by the positive control primer pair is 200 base pairs.

Length of PCR	105	200	105	80	75	80	85
product							
5'-primer ¹	164	340	440	45	45	43	36
•	5'-CAC3'	^{5'} -Agg ^{3'}	^{5'} -TTA3'	^{5'} -Tgg ^{3'}	^{5'} -Tgg ^{3'}	^{5'} -Tgg ^{3'}	5'-TAC3'
							36
							^{5'} -TAT ^{3'}
3'-primer ²	231	2 nd I	507	59	58	57	47
	⁵ '-TgC ³ '	^{5'} -AAA ^{3'}	⁵ '-TTg ³ '	^{5'} -CTC ^{3'}	^{5'} -ggC ^{3'}	5'-CTC ^{3'}	5'-ACA3'
							48
							^{5'} -gCA ^{3'}
							48
							^{5'} -gCC ^{3'}
							52
							^{5'} -TgT ^{3'}
A*	+	+	+				
B*	+	+	+				
C*	+	+	+				
DRB1				+	+		
DRB3				+	+		
DRB5				+			
DQB1					+		
DPB1						+	
DQA1	tion for I			1 41			+

¹The nucleotide position for HLA class I genes and the codon for HLA class Il genes, in the 2nd or 3rd exon, matching the specificity-determining 3'-end of the primer is given. Nucleotide and codon numbering as on the www.ebi.ac.uk/imgt/hla web site. The sequence of the 3 terminal nucleotides of the primer is given.

²The nucleotide position for HLA class I genes and the codon for HLA class Il genes, in the 2nd or 3rd exon or the 2nd intron, matching the specificitydetermining 3'-end of the primer is given in the anti-sense direction. Nucleotide and codon numbering as on the www.ebi.ac.uk/imgt/hla web site. The sequence of the 3 terminal nucleotides of the primer is given.

•LERUP SSP*

HLA-C*04:09N Product Insert 101.861-12 – including *Taq* polymerase, IFU-01 101.861-12u – without *Taq* polymerase, IFU-02

Page 3 of 8
Visit https://labproducts.caredx.com for "Instructions for Use" (IFU)

Lot No.: **4L4** Lot-specific information

PRODUCT DESCRIPTION

HLA-C*04:09N SSP subtyping

CONTENT

The primer set contains 5'- and 3'-primers for identifying the HLA-C*04:09N allele.

PLATE LAYOUT

Each test consists of 3 PCR reactions in an 8 well PCR plate. Wells 4 to 8 are empty.

1 2 NC empty empty empty empty empty

The 8 well cut PCR plate is marked with the Lot No. '4L4' in silver/gray ink.

Well No. 1 is marked with the Lot No. '4L4'.

Wells 1 to 2 – HLA-C*04:09N high resolution primers.

Well 3 – Negative Control (NC).

A faint row of numbers is seen between wells 1 and 2 or wells 7 and 8 of the PCR trays. These stem from the manufacture of the trays, and should be disregarded. The PCR plates are heat-sealed with a PCR-compatible foil.

Please note: When removing each 8 well PCR plate, make sure that the remaining plates stay sealed. Use a scalpel or a similar instrument to carefully cut the foil between the plates.

INTERPRETATION

Due to the sharing of sequence motifs between HLA-C alleles, non-HLA-C*04:09N alleles will be amplified by primer mix 2. For further details see Specificity Table.

UNIQUELY IDENTIFIED ALLELES

The HLA-C*04:09N allele will give rise to a unique amplification pattern by the primers in the HLA-C*04:09N kit^{1,2}.

¹HLA-C alleles listed on the IMGT/HLA web page 2020-April-20, release 3.40.0, www.ebi.ac.uk/imgt/hla.

²Alleles that have been deleted from or renamed in the official WHO HLA Nomenclature up to and including the last IMGT/HLA database release can be retrieved from web page http://hla.alleles.org/alleles/deleted.html.

((

Visit https://labproducts.caredx.com for "Instructions for Use" (IFU)

Lot No.: 4L4

Lot-specific information SPECIFICITY TABLE

HLA-C*04:09N SSP subtyping

Specificities and sizes of the PCR products of the 2+1 primer mixes used for **HLA-C*04:09N SSP subtyping**

Primer Mix	Size of spec. PCR product ¹	Size of control band ²	Amplified HLA-C alleles ³
1	220 bp	800 bp	*04:09N
2	260 bp, 360 bp	1070 bp	*01:177, 01:186, 03:231, 03:248, 03:308, 04:01:01:01-04:01:01:29, 04:01:01:31-04:01:120, 04:03:01:01-04:20, 04:23-04:212, 04:214-04:225N, 04:227-04:350N, 04:352-04:413, 05:78:01-05:78:02, 05:141, 06:120, 08:163, 08:183, 12:194, 12:269, 16:34, 16:62, 16:150
3 ⁴	-	-	Negative Control

¹Alleles are assigned by the presence of specific PCR product(s). However, the sizes of the specific PCR products may be helpful in the interpretation of HLA-C*04:09N SSP typings.

Nonspecific amplifications, i.e. a ladder or a smear of bands, may sometimes be seen. GC-rich primers have a higher tendency of giving rise to nonspecific amplifications than other primers.

PCR fragments longer than the control bands may sometimes be observed. Such bands should be disregarded and do not influence the interpretation of the SSP typings.

PCR fragments migrating faster than the control bands, but slower than a 400 bp fragment may be seen in some gel read-outs. Such bands can be disregarded and do not influence the interpretation of the SSP typings.

Some primers may give rise to primer oligomer artifacts. Sometimes this phenomenon is an inherit feature of the primer pair(s) of a primer mix. More often it is due to other factors such as too low amount of DNA in the PCR reactions, taking too long time in setting up the PCR reactions, working at elevated room temperature or using thermal cyclers that are not pre-heated.

²The internal positive control primer pairs amplify segments of the human growth hormone gene. The internal positive control bands are 1070 or 800 base pairs respectively, well distribution as outlined in the table. Well number 1 contains the shorter, 800 bp, internal positive control band. The well distribution of the internal controls can help in orientation of the kit on gel photo, as well as allow for kit identification. In the presence of a specific amplification the intensity of the control band often decreases.

³For several HLA Class I alleles 1st and/or 4th exon(s) and beyond, as well as intron nucleotide sequences, are not available. In these instances it is not known whether some of the primers of the SSP sets are completely matched with the target sequences or not. Assumption is made that unknown sequences in these regions are conserved within allelic groups.

⁴Primer mix 3 contains a negative control, which will amplify the majority of HLA amplicons as well as the amplicons generated by the control primer pairs matching the human growth hormone gene. HLA-specific PCR product sizes range from 75 to 200 base pairs and the PCR product generated by the HGH positive control primer pair is 200 base pairs.

"Instructions for Use" (IFU)

Lot No.: 4L4

Lot-specific information

PRIMER SPECIFICATION

Well No.	1	2
Length of spec.	220	260
PCR product		360
Length of int.	800	1070
pos. control ¹		
5'-primer(s) ²	1018	28
	^{5'} -gTg ^{3'}	^{5'} -TCA ^{3'}
3'-primer(s) ³	1092	112
	^{5'} -TTA ^{3'}	5' -CCA 3'
		118
		^{5'} -gCT ^{3'}
		218
		^{5'} -gCT ^{3'}
Well No.	1	2

¹The internal positive control primer pairs amplify segments of the human growth hormone gene. The internal positive control bands are 1070 or 800 base pairs respectively, well distribution as outlined in the table. Well number 1 contains the shorter, 800 bp, internal positive control band. The well distribution of the internal controls can help in orientation of the kit on gel photo, as well as allow for kit identification. In the presence of a specific amplification the intensity of the control band often decreases.

²The nucleotide position matching the specificity-determining 3'-end of the primer is given. Nucleotide numbering as on the www.ebi.ac.uk/imgt/hla web site. The sequence of the 3 terminal nucleotides of the primer is given.

³The nucleotide position matching the specificity-determining 3'-end of the primer is given in the anti-sense direction. Nucleotide numbering as on the www.ebi.ac.uk/imgt/hla web site. The sequence of the 3 terminal nucleotides of the primer is given.

HLA-C*04:09N

Lot No.: **4L4 Lot-specific information**

		INE VALI				
	H	LA-C*04:0	9N SSI	2 kit ²		
					<u> </u>	ell
					1	2
				Prod No.:	201783801	201783802
	IHV	C cell line ¹		C*		
1	9001		*07:02		-	-
2	9280	LK707	*07:01	*15:05	-	-
3	9011	E4181324	*12:02		-	-
4	9275	GU373	*03:04	*04:01	-	+
5	9009	KAS011	*06:02		-	-
6	9353	SM	*03:04	*07:02	-	-
7	9020	QBL	*05:01		-	-
8	9007	DEM	*04:01		-	+
9	9026	YAR	*12:03		-	-
10	9107	LKT3	*01:02		-	-
11		PITOUT	*16:01		-	-
12	9052	DBB	*06:02		-	-
13	9004	JESTHOM	*01:02		-	-
14		OLGA	*01:02	*03:04	-	-
15	9075		*03:04		-	-
16		SWEIG007	*02:02		-	-
17		CTM3953540	*03:03	*07:01	-	-
18		32367	*01:02	*07:05		-
19		BM16	*07:01	07.00		
20		SLE005	*03:04		-	H
21		AMALA	*03:03	1	H	-
22		KOSE	*12:03		H	-
	9124		*01:02	*15,00	-	-
23 24		JBUSH	*12:03	*15:02	-	-
_					-	Ē
25		IBW9	*08:02		-	-
26		WT49	*07:18	*45.00	-	-
27		CH1007	*07:04	*15:29	-	-
28		BEL5GB	*05:01	*16:01	-	-
29		MOU	*16:01		-	-
30	9021		*17:01		-	-
31		DUCAF	*05:01		-	-
32		HAG	*17:03		-	-
33		MT14B	*03:04		-	-
34	9104		*12:03		-	-
35	9302	SSTO	*05:01		-	-
36	9024	KT17	*03:03	*04:01	-	+
37	9065	HHKB	*07:02		-	-
38	9099	LZL	*03:03		-	-
39	9315	CML	*02:02	*07:01	-	-
40	9134	WHONP199	*01:02	*06:02	-	-
41	9055	H0301	*08:02		-	-
42		TAB089	*01:02		-	-
43		T7526	*01:02	*08:01	-	-
44	9057		*12:03		-	-
45		SHJO	*06:02	*17:01	-	-
46		SCHU	*07:02		-	-
47		TUBO	*07:04	*15:02	-	-
48		TER-ND	*04:01	*16:01	-	+

OLERUP SSP

HLA-C*04:09N **Product Insert** Page 7 of 8 **101.861-12 – including** *Taq* **polymerase**, IFU-01 Visit https://labproducts.caredx.com for **101.861-12u – without** *Taq* **polymerase**, IFU-02 "Instructions for Use" (IFU)

Lot No.: 4L4 **Lot-specific information**

¹The provided cell line HLA specificities are retrieved from the http://www.ihwg.org/hla web site. The specificity

of an individual cell line may thus be subject to change.

²The specificity of each primer solution in the kit has been tested against 48 well characterized cell line DNAs and where applicable, additional cell line DNAs.

OLERUP SSP*

HLA-C*04:09N Product Insert Page 8 of 8
101.861-12 – including *Taq* polymerase, IFU-01 Visit https://labproducts.caredx.com for 101.861-12u – without *Taq* polymerase, IFU-02 "Instructions for Use" (IFU)

Lot No.: **4L4** Lot-specific information

ADDRESSES:

Manufacturer:

CareDx AB, Franzengatan 5, SE-112 51 Stockholm, Sweden.

Tel: +46-8-508 939 00 Fax: +46-8-717 88 18

E-mail: orders-se@caredx.com

Web page: https://labproducts.caredx.com/

Distributed by:

CareDx GmbH, Löwengasse 47 / 6, AT-1030 Vienna, Austria.

Tel: +43-1-710 15 00 **Fax:** +43-1-710 15 00 10 **E-mail:** orders-at@caredx.com

Web page: https://labproducts.caredx.com/

CareDx Lab Solutions Inc., 901 S. Bolmar St., Suite R, West Chester, PA 19382

Tel: 1-877-653-78171 **Fax:** 610-344-7989

E-mail: orders-us@caredx.com

Web page: https://labproducts.caredx.com/

For information on CareDx distributors worldwide, contact **CareDx GmbH**.