Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information

Olerup SSP[®] HLA-C*15 Add-on

Product number:	101.865-12 – including <i>Taq</i> polymerase 101.865-12u – without <i>Taq</i> polymerase
Lot number:	5E7
Expiry date:	2019-05-01
Number of tests:	12
Number of wells per test:	3+1
Storage - pre-aliquoted primers:	dark at -20°C
- PCR Master Mix:	-20°C
 Adhesive PCR seals 	RT
- Product Insert	RT

This Product Description is only valid for Lot No. 5E7.

Complete product documentation consists of generic Instructions for Use (IFU), lot specific Product Insert, Worksheet and Certificate.

CHANGES COMPARED TO THE PREVIOUS OLERUP SSP[®] HLA-C*15 ADD-ON LOT (91X)

The HLA-C*15 Add-on specificity and interpretation tables have been updated for the HLA-C alleles described since the previous *Olerup* SSP[®] HLA-C*15 Add-on lot was made **(Lot No. 91X)**. The kit design is based on IMGT/HLA database 3.25.0.

As of lot series V, the Specificity Table is included in the lot-specific Product Insert, and the Interpretation Table is included in the Worksheet.

The primers of the wells detailed below have been exchanged, added or modified compared to the previous lot.

Well	5'-primer	3'-primer	rationale
4	-	-	Updated Negative Control.

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information

Well **4** contains <u>Negative Control primer pairs</u>, that will amplify more than 95% of the *Olerup* SSP[®] HLA Class I, DRB, DQB1, DPB1 and DQA1 amplicons as well as all the amplicons generated by the control primer pairs matching the human growth hormone gene.

HLA-specific PCR product sizes range from 75 to 200 base pairs. The PCR product generated by the positive control primer pair is 430 base pairs.

Length of PCR	105	200	105	80	75	80	85
product							
5'-primer ¹	164	340	440	45	45	43	36
	^{5'} -CAC ^{3'}	^{5'} -Agg ^{3'}	^{5′} -TTA3′	^{5′} -Tgg³′	⁵'-Tgg³'	^{5′} -Tgg ^{3′}	^{5'} -TAC ^{3'}
							36
							^{5'} -TAT ^{3'}
3'-primer ²	231	2 nd I	507	59	58	57	47
-	^{5'} -TgC ^{3'}	^{5'} -AAA ^{3'}	^{5'} -TTg ^{3'}	^{5'} -CTC ^{3'}	^{5'} -ggC ^{3'}	^{5'} -CTC ^{3'}	^{5'} -ACA ^{3'}
							48
							^{5'} -gCA ^{3'}
							48
							^{5'} -gCC ^{3'}
							52
							^{5′} -TgT ^{3′}
A*	+	+	+				
B*	+	+	+				
C*	+	+	÷				
DRB1				+	+		
DRB3				+	+		
DRB5				+			
DQB1					+		
DPB1						+	
DQA1							+

¹The nucleotide position for HLA class I genes and the codon for HLA class II genes, in the 2nd or 3rd exon, matching the specificity-determining 3'-end of the primer is given. Nucleotide and codonnumbering as on the <u>www.ebi.ac.uk/imgt/hla</u> web site. The sequence of the 3 terminal nucleotides of the primer is given.

²The nucleotide position for HLA class I genes and the codon for HLA class II genes, in the 2nd or 3rd exon or the 2nd intron, matching the specificitydetermining 3'-end of the primer is given in the anti-sense direction. Nucleotide and codon numbering as on the <u>www.ebi.ac.uk/imgt/hla</u> web site. The sequence of the 3 terminal nucleotides of the primer is given.

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information PRODUCT DESCRIPTION

HLA-C*15 Add-on SSP subtyping

CONTENT

The primer set contains 5'- and 3'-primers for distinguishing the HLA-C*15:29 from the C*15:05 alleles and the C*15:87 from the C*15:02 alleles.

PLATE LAYOUT

Each test consists of 4 PCR reactions in an 8 well cut PCR plate. Wells 5 to 8 are empty.

1 2 3 NC empty empty empty empty

The 8 well cut PCR plate is marked with the Lot No. '5E7' in silver/gray ink. Well No. 1 is marked with the Lot No. '5E7'.

Wells 1 to 3 – HLA-C*15 Add-on high resolution primers.

Well 4 – Negative Control (NC).

A faint row of numbers is seen between wells 1 and 2 or wells 7 and 8 of the PCR trays. These stem from the manufacture of the trays, and should be disregarded. The PCR plates are heat-sealed with a PCR-compatible foil.

Please note: When removing each 8 well PCR plate, make sure that the remaining plates stay sealed. Use a scalpel or a similar instrument to carefully cut the foil between the plates.

INTERPRETATION

The interpretation of HLA-C*15 Add-on SSP subtypings will be influenced by a few other C*15 and HLA-C alleles. In addition, two HLA-B alleles will be amplified by primer mix 1.

For further details see Specificity Table.

UNIQUELY IDENTIFIED ALLELES

The HLA-C^{*} C^{*}15:02, 15:05, 15:29, and C^{*}15:87 alleles give different patterns in the HLA-C^{*}15 Add-on kit^{1,2}.

The HLA-C*15 Add-on kit cannot distinguish the silent mutations in the C*15:02:01:01-15:02:26 or the C*15:05:01-15:05:12 alleles.

¹Based on HLA-C alleles listed on the IMGT/HLA web page 2016-July-14, release 3.25.0, <u>www.ebi.ac.uk/imgt/hla</u>.

²Alleles that have been deleted from or renamed in the official WHO HLA Nomenclature up to and including the last IMGT/HLA database release can be retrieved from web page http://hla.alleles.org/alleles/deleted.html.

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information SPECIFICITY TABLE

HLA-C*15 Add-on SSP subtyping

Specificities and sizes of the PCR products of the 3+1 primer mixes used for HLA-C*15 Add-on SSP subtyping

Primer Mix	Size of spec. PCR product ¹	Size of control band ²	Amplified HLA- C*15:02/15:05/15:29/15:87 alleles ³	Other amplified HLA-C alleles ⁴
15	100 bp	800 bp	*15:05:01-15:05:12, 15:29	*15:22-15:23, 15:36, 15:46, 15:54, 15:59, 15:69-15:70, 15:90, 15:104, 15:108, 15:110-15:111, 15:115N- 15:117, 15:125, 04:108, B*07:246, B*35:03:11, B*35:205
2	545 bp	1070 bp	*15:29, 15:87	*01:85, 08:22, 08:56
3	315 bp	800 bp	*15:02:01-15:02:26	*15:03, 15:07-15:08, 15:10:01-15:13, 15:15- 15:18, 15:21, 15:26, 15:28, 15:31-15:35, 15:37-15:39, 15:41-15:45, 15:47-15:53, 15:56-15:58, 15:60, 15:62- 15:63, 15:67-15:68, 15:71- 15:75, 15:78-15:89, 15:91- 15:101, 15:103, 15:106, 15:109, 15:112-15:114, 15:118-15:124, 15:127, 15:129, 02:06:01-02:06:02, 02:47, 03:19, 03:102, 03:307, 04:178, 05:132, 06:168, 07:289, 12:15
4 ⁶	-	-	Negative Control	

¹Alleles are assigned by the presence of specific PCR product(s). However, the sizes of the specific PCR products may be helpful in the interpretation of HLA-C*15 Add-on SSP typings.

Nonspecific amplifications, i.e. a ladder or a smear of bands, may sometimes be seen. GC-rich primers have a higher tendency of giving rise to nonspecific amplifications than other primers.

PCR fragments longer than the control bands may sometimes be observed. Such bands should be disregarded and do not influence the interpretation of the SSP typings.

PCR fragments migrating faster than the control bands, but slower than a 400 bp fragment may be seen in some gel read-outs. Such bands can be disregarded and do not influence the interpretation of the SSP typings.

Some primers may give rise to primer oligomer artifacts. Sometimes this phenomenon is an inherit feature of the primer pair(s) of a primer mix. More often it is due to other factors such as too low amount of DNA in the PCR reactions, taking too long time in setting up the PCR reactions, working at elevated room temperature or using thermal cyclers that are not pre-heated.

²The internal positive control primer pairs amplify segments of the human growth hormone gene. The internal positive control bands are 1070 or 800 base pairs respectively, well distribution as outlined in the table. Well number 1 contains the shorter, 800 bp, internal positive control band. The well distribution of the internal controls can help in orientation of the kit on gel photo, as well as allow for kit identification. In the presence of a specific amplification the intensity of the control band often decreases.

Product Insert

101.865-12 – including *Taq* polymerase, IFU-01 101.865-12u – without *Taq* polymerase, IFU-02

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information

³For several HLA Class I alleles 1st and/or 4th exon(s) and beyond, as well as intron nucleotide sequences, are not available. In these instances it is not known whether some of the primers of the SSP sets are completely matched with the target sequences or not. Assumption is made that unknown sequences in these regions are conserved within allelic groups.

⁴The interpretation of HLA-C*15 Add-on SSP subtypings will be influenced by a few other C*15 and HLA-C alleles. In addition, two HLA-B alleles will be amplified by primer mix 1.

⁵HLA-specific PCR products shorter than 125 base pairs have a lower intensity and are less sharp than longer PCR products.

⁶Primer mix 4 contains a negative control, which will amplify more than 95% of HLA amplicons as well as the amplicons generated by the control primer pairs matching the human growth hormone gene. HLA-specific PCR product sizes range from 75 to 200 base pairs and the PCR product generated by the HGH positive control primer pair is 430 base pairs.

Visit www.olerup-ssp.com for

"Instructions for Use" (IFU)

101.865-12 – including *Taq* polymerase, IFU-01 101.865-12u – without *Taq* polymerase, IFU-02

Lot No.: 5E7

Lot-specific information PRIMER SPECIFICATION

Well No.	1	2	3
Length of spec.	100	545	315
PCR product			
Length of int.	800	1070	800
pos. control ¹			
5'-primer(s) ²	420	972	420
	^{5'} -TTC ^{3'}	^{5'} -CTA ^{3'}	^{5'} -TTA ^{3'}
			420
			^{5'} -TTA ^{3'}
3'-primer(s) ³	477	1034	3 rd I
	^{5'} -gCg ^{3'}	^{5'} -AgT ^{3'}	^{5'} -CTC ^{3'}
Well No.	1	2	3

¹The internal positive control primer pairs amplify segments of the human growth hormone gene. The internal positive control bands are 1070 or 800 base pairs respectively, well distribution as outlined in the table. Well number 1 contains the shorter, 800 bp, internal positive control band. The well distribution of the internal controls can help in orientation of the kit on gel photo, as well as allow for kit identification. In the presence of a specific amplification the intensity of the control band often decreases.

²The nucleotide position matching the specificity-determining 3'-end of the primer is given. Nucleotide numbering as on the <u>www.ebi.ac.uk/imgt/hla</u> web site. The sequence of the 3 terminal nucleotides of the primer is given.

³The nucleotide position matching the specificity-determining 3'-end of the primer is given in the anti-sense direction. Nucleotide numbering as on the <u>www.ebi.ac.uk/imgt/hla</u> web site. The sequence of the 3 terminal nucleotides of the primer is given.

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

101.865-12 – including *Taq* polymerase, IFU-01 101.865-12u – without *Taq* polymerase, IFU-02

Lot No.: 5E7

Lot-specific information

Lot	No.: 5	E/		Lot-sp	eci	fic	info	rm
С		LINE VAI			-	E	Т	
	н	LA-C*15 A	dd-on S	SSP kiť	2			
						Nel	I	
					1	2	3	
				Lot No.:	201296601	201296602	201547603	
	IHM	/C cell line ¹		C*				
1	9001	SA	*07:02		-	-	-	
2	9280	LK707	*07:01	*15:05	+	-	-	
3	9011	E4181324	*12:02		-	-	-	
4	9275	GU373	*03:04	*04:01	-	-	-	
5		KAS011	*06:02		-	-	-	
6	9353		*03:04	*07:02	-	-	-	
7	9020		*05:01		-	-	-	
8	9025	DEU	*04:01		-	-	-	
9		YAR	*12:03		-	-	-	
10	9107	LKT3	*01:02		-	-	-	
11	9051	PITOUT	*16:01		-	-	-	
12	9052	DBB	*06:02		-	-	-	
13	9004	JESTHOM	*01:02		-	-	-	
14	9071	OLGA	*01:02	*03:04	-	-	-	
15	9075	DKB	*03:04		-	-	-	
16	9037	SWEIG007	*02:02		-	-	-	
17	9282	CTM3953540	*03:03	*07:01	-	-	-	
18	9257	32367	*01:02	*07:05	-	-	-	
19		BM16	*07:01		-	-	-	
20		SLE005	*03:04		-	-	-	
21		AMALA	*03:03		-	-	-	
22		KOSE	*12:03		-	-	-	
23	9124		*01:02	*15:02	-	-	+	
24		JBUSH	*12:03		-	-	-	
25		IBW9	*08:02		-	-	-	
26		WT49	*07:01		-	-	-	
27		CH1007	*07:04	*15:29	+	+	-	
28		BEL5GB	*05:01	*16:01	-		-	
20 29	9320		*16:01	10.01		-	-	
29 30	9050		*17:01		1_	-	-	
30 31		DUCAF	*05:01		-	-		
31 32		HAG	*17:01	*17:03			-	
		MT14B	*03:04	17.03	-	-	-	
33 24					-	-	-	
34 25	9104		*12:03		-	-	-	
35 26		SSTO	*05:01	*04.04	-	-	-	
36		KT17	*03:03	*04:01	-	-	-	
37		HHKB	*07:02		-	-	-	
38	9099		*03:03	*07-04	-	-	-	
39 40	9315		*02:02	*07:01	-	-	-	
40		WHONP199	*01:02	*06:02	-	-	-	
41		H0301	*08:02		-	-	-	
42		TAB089	*01:02	105 5	-	-	-	
43		T7526	*01:02	*08:01	-	-	-	
44	9057		*12:03		-	-	-	
45		SHJO	*06:02	*17:01	-	-	-	
46		SCHU	*07:02		-	-	-	
47		TUBO	*07:04	*15:02	-	-	+	
48	9303	TER-ND	*04:01	*16:01	-	-	-	

Product Insert

101.865-12 – including *Taq* polymerase, IFU-01 101.865-12u – without *Taq* polymerase, IFU-02

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information

¹The provided cell line HLA specificities are retrieved from the <u>http://www.ihwg.org/hla</u> web site. The specificity of an individual cell line may thus be subject to change.

²The specificity of each primer solution in the kit has been tested against 48 well characterized cell line DNAs and where applicable, additional cell line DNAs.

In primer solution 3 one 5'-primer was not possible to test.

Lot No.: 5E7

Lot-specific information

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7

Lot-specific information

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Visit <u>www.olerup-ssp.com</u> for "Instructions for Use" (IFU)

Lot No.: 5E7 ADDRESSES: Lot-specific information

Manufacturer: Olerup SSP AB, Franzengatan 5, SE-112 51 Stockholm, Sweden. *Tel:* +46-8-717 88 27 *Fax:* +46-8-717 88 18 *E-mail:* info-ssp@olerup.com *Web page:* http://www.olerup-ssp.com

Distributed by: Olerup GmbH, Löwengasse 47 / 6, AT-1030 Vienna, Austria. *Tel:* +43-1-710 15 00 *Fax:* +43-1-710 15 00 10 *E-mail:* support-at@olerup.com *Web page:* http://www.olerup.com

Olerup Inc., 901 S. Bolmar St., Suite R, West Chester, PA 19382 *Tel:* 1-877-OLERUP1 *Fax:* 610-344-7989 *E-mail:* info.us@olerup.com *Web page:* http://www.olerup.com

For information on Olerup SSP distributors worldwide, contact Olerup GmbH.