101.612-12 – including *Taq* polymerase 101.612-12u – without *Taq* polymerase

Lot No.: **60R**

Olerup SSP® HLA-C*04

Product number: 101.612-12 – including *Taq* polymerase

101.612-12u – without *Taq* polymerase

Lot number: 60R

Expiry date: 2015-June-01

Number of tests: 12 Number of wells per test: 48

Changes compared to the previous HLA-C*04 Lot (79N):

| Well | 5'-primer | 3'-primer | rationale |
|------|-----------|-----------|---|
| 9 | - | - | Exchanged positive control primer pair. |
| 14 | - | Removed | 3'-primer removed for improved specificity of |
| | | | amplification and decreased tendencies of |
| | | | primer oligomer formation. |
| 16 | Added | - | 5'-primer added for the C*04:123N allele. |
| 27 | - | Added | 3'-primer added for the C*04:120 allele. |
| 29 | Added | Added | Primer pair added for the C*04:82 allele. |
| 34 | - | Added | 3'-primer added for the C*04:120 allele. |
| 35 | - | Added | 3'-primer added for the C*04:115N allele. |
| 38 | Added | - | 5'-primer added for the C*04:123N allele. |
| 40 | - | Added | 3'-primer added for the C*04:64:02 allele. |
| 42 | Exchanged | Exchanged | Exchanged primer pair for the C*04:82 |
| | | | allele. |
| 43 | - | Added | 3'-primer added for the C*04:115N allele. |
| 44 | Added | - | 5'-primer added for the C*04:108 allele. |
| 47 | Moved, | Moved, | Primer pairs moved to well 48, new primer |
| | added | added | pair for the C*04:114 allele. |
| 48 | New | New | Primer pair from well 47. |

THE NUMBER OF WELLS has been increased from 47 to 48.

ALLELE COVERAGE:

C*04:01 to C*04:127, i.e. all the currently recognized HLA-C*04 alleles, will be amplified by the primers in the HLA-C*04 kit ¹; www.ebi.ac.uk/imgt/hla, 2012-July-12, release 3.9.0.

The HLA-C*04 kit enables separation of the confirmed HLA-C*04 alleles as listed in the IMGT/HLA database. An HLA allele is listed as confirmed by IMGT/HLA if it has been sequenced by more than a single laboratory or from multiple sources.

The HLA-C*04 kit also enables identification of polymorphisms in exons outside of the region encoding the peptide binding domain and of null and alternatively expressed alleles.

December 2012
Rev. No.: 00

101.612-12 – including *Taq* polymerase 101.612-12u – without *Taq* polymerase

Lot No.: **60R**

The HLA-C*04 primer set cannot distinguish the following silent mutations: the C*04:01:01:01-04:01:09, 04:01:11-04:01:22, 04:01:24-04:01:27 and 04:01:29-04:01:41 alleles, the C*04:01:10 and 04:01:23 alleles, the C*04:04:01-04:04:02 alleles, the C*04:15:01 and 04:15:03 alleles, the C*04:64:01-04:64:02 alleles and the C*04:94:01-04:94:02 alleles.

¹The C*04:15:02, C*04:44 and C*04:47 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 20.

The C*04:23, C*04:38 and C*04:39 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 21.

The C*04:25, C*04:40 and C*04:41 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 23.

The C*04:46 and C*04:50 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 27.

The C*04:48 and C*04:75 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 28.

The C*04:54 and C*04:105N alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 41.

The C*04:56 and C*04:64:01-04:64:02 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 40.

The C*04:59Q and C*04:78 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 32.

The C*04:61 and C*04:96 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 34.

The C*04:62 and C*04:76 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 35.

The C*04:65 and C*04:72 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 33.

The C*04:67 and C*04:93N alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 46.

The C*04:81 and C*04:88N alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 43.

The C*04:84 and C*04:106 alleles can be distinguished by the different sizes of the specific PCR products generated by primer mix 48.

²The HLA-C*04 primer set cannot separate the C*04:03 and 02:49 alleles or the C*04:107 and the C*02:12 and 02:55 alleles. These alleles can be distinguished by the HLA-C low resolution kit and/or the HLA-C*02 high resolution kit.

RESOLUTION IN HLA-C*04 HOMO- AND HETEROZYGOTES: Good.

INFLUENCE ON THE INTERPRETATION OF HLA-C*04 SUBTYPINGS BY NON-HLA-C*04 ALLELES:

None frequently occurring.

MODIFICATIONS MADE DUE TO COMMENTS FROM CUSTOMERS:

No comments received.